

What is Claimed is:

1. An E1-deficient adenovirus comprising a YB-1 encoding DNA sequence.
2. An E1-deficient adenovirus as in claim 1, wherein said adenovirus is an E1A-deficient adenovirus.
3. A nucleic acid encoding an E1-deficient adenovirus comprising a YB-1 encoding DNA sequence.
4. A nucleic acid as in claim 3, wherein said adenovirus is an E1A-deficient adenovirus.
5. A pharmaceutical composition comprising an E1-deficient adenovirus comprising a YB-1 encoding DNA sequence and at least one pharmaceutically acceptable carrier.
6. A pharmaceutical composition as in claim 5, wherein said adenovirus is an E1A-deficient adenovirus.
7. A method for treatment of tumors, comprising administering to a patient in need thereof an anti-tumor effective amount of a medicament comprising an E1-deficient adenovirus comprising a YB-1 encoding DNA sequence.
8. A method as in claim 7, wherein said adenovirus is an E1A-deficient adenovirus.
9. A method as in claim 7, wherein said medicament is carried on at least one pharmaceutically acceptable carrier.

10. A method as in claim 7, wherein said method includes at least one of:
administration of substances which damage tumor cells;
surgical tumor excision;
radiation therapy;
chemotherapy;
hyperthermia; and
gene therapy.
11. A method as in claim 10, wherein said substances which damage tumor cells are selected from the group consisting of cytostatic agents and ribozymes.
12. A method according to claim 7, wherein said tumor cells are tumor cells which exhibit YB-1 in the nucleus.
13. A method for E1-independent replication of a replication-defective adenovirus comprising
administering to a cell a recombinant adenovirus carrying a YB-1 encoding DNA sequence,
inducing expression of YB-1 in said cell,
causing said adenovirus to replicate in said cell in the presence of YB-1.
14. A method as in claim 13, wherein the adenovirus is a E1-deficient adenovirus.
15. A method as in claim 13, wherein the adenovirus is a E1A-deficient adenovirus.
16. A method as in claim 13, wherein YB-1 is expressed in the nucleus of said cell.
17. A method for E1-independent replication of a replication-defective adenovirus comprising
administering to a cell a replication-defective adenovirus, whereby YB-1 is present in the nucleus of the cell,

11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66																																		

19. A method as in claim 17, wherein the adenovirus is a E1-deficient adenovirus.

- 13 -